

Subject: Treatment Guidelines - ALS  
**Severe Pre-Eclampsia/ Eclampsia**

Associated Policies:

---

I. Priorities


- A. ABC's
- B. Identify signs of shock.
- C. If pregnant, determine stage (trimester) of pregnancy.
- D. Determine the degree of physiologic distress, estimate amount of blood loss.
- E. Rapid transport, if unconscious or conscious with progressive symptoms.

II. Severe Pre-Eclampsia/ Eclampsia

Third trimester pregnancy with hypertension (BP systolic > 160, diastolic > 110), mental status changes, visual disturbances, peripheral edema (pre-eclampsia), seizures and/or coma (eclampsia).

- A. Ensure a patent airway
- B. Administer Oxygen per Policy # 6030.
- C. Position mother on her left side.
- D. Transport quickly and maintain a quiet environment.
- E. IV access TKO while en route.
- F. Determine blood glucose. Treat hypoglycemia per Policy #6515.
- G. Administer Mag Sulfate 10% per Policy #5428 in the presence of ALOC or seizures in the third trimester pregnancy.
- H. Be prepared to treat seizures per Policy # 6516 if Mag Sulfate 10% is delayed.
- I. Transport to appropriate facility.

Approved:  Date: 4/24/20

Approved as to Form:  Date: 4/24/20